

## Prevalence and In-vitro Evaluation of Probiotic Properties of *Lactobacillus delbrueckii* and *Lactobacillus plantarum* Isolated from Yoghurt in Chittagong Division, Bangladesh

Abu Sayeed Mohammad Mohamud<sup>1\*</sup>, Mobarak Chowdhury<sup>2</sup>, Rezaul Karim<sup>4</sup>, Saiful Islam<sup>1</sup>, Md. Saidur Rahman<sup>1</sup>, Milton Halder<sup>1</sup>, Khorshed Alam<sup>3</sup>, Habibur Rahman Bhuiyan<sup>1</sup> and Faridul Islam<sup>1</sup>

<sup>1</sup>Bangladesh Council of Scientific and Industrial Research Laboratories, Chittagong, Bangladesh.

<sup>2</sup>Shahjalal University of Science and Technology, Sylhet, Bangladesh.

<sup>3</sup>Bangladesh Standards and Testing Institution.

<sup>4</sup>Institute of Food Science and Technology, Bangladesh Council of Scientific and Industrial Research.

(Received: 14 April 2014; accepted: 28 May 2014)

A total of five different Lactic acid bacterial species were identified from yoghurt samples collected from Chittagong region of Bangladesh. Among the isolates, the prevalence of *Lactobacillus delbrueckii* and *Lactobacillus plantarum* were 111 and 114cfu/ml respectively. After identification and isolation, we have optimized the suitable growth of isolates against pH, temperature and various concentration of NaCl. The results showed that better growth of organisms was observed in the presence of 3-6% NaCl. These species can tolerate up to 3% bile salt but the best tolerance found at 1% bile salt. To evaluate the antimicrobial properties of identified species, growth inhibition test has been done against some selected pathogens. It showed that Lactobacilli inhibited the growth of all pathogenic bacterial species. Inhibition by isolates and further treated with papain indicate that our isolates inhibited the growth of pathogenic microorganisms by producing Bacteriocins like substances. The selected species were subjected to Antibiotic Susceptibility testing to observe their features of carrying Antibiotic resistance genes. Among the 21 Antibiotics 17 antibiotics found resistant to *Lactobacillus delbrueckii*, on the other hand, *Lactobacillus plantarum* were found resistant to 4 antibiotics.

**Key words:** Lactic Acid Bacteria, Probiotic, Lactobacillus, Prevalence, Yoghurt.

Yoghurt is a widely enjoyed dairy product that is essentially an altered form of milk containing waste products from fermentation. The lactic acid that is produced from the fermentation of lactose contributes to the sour taste of yogurt by decreasing pH and allows for the characteristic texture by acting on the milk proteins<sup>18</sup>. The acids created by the friendly bacteria fermenting the milk

help to curb the growth of unfriendly bacteria in the yogurt. Yogurt has been continually studied for its health benefits, particularly from the addition of probiotics. Current research has been investigating how to improve yogurt both in terms of its potential as a healthy food and as an appetizing product that appeals to the general population. The probiotic efficacy of lactic acid bacteria isolated from traditional Ethiopian fermented foods<sup>6</sup>. The prevalence of lactic acid bacteria (LAB) was studied in Senegalese local food products was determined to be 109 CFU/g in millet flour and milk products, and 103 CFU/g in seafood products<sup>6</sup>. Critical evaluation of probiotic

\* To whom all correspondence should be addressed.  
Mobile: +8801746700196; Fax: +031682505;  
E-mail: sayeedmahmud@bcsir.gov.bd

activity of lactic acid bacteria and their effects was studied<sup>10</sup>. Potential activities of *Lactobacillus* exopolysaccharides was studied on immunomodulatory and antioxidant<sup>1</sup>. The probiotic properties of *Lactobacillus* isolates originating from porcine intestine and feces was also observed<sup>12</sup>. The cancer preventing attributes of probiotics was investigated<sup>11</sup>. In Bangladesh, yoghurt is perhaps the oldest fermented milk product known and consumed by large sectors of the population as a part of their daily diet. In most of the areas of Bangladesh, different types of traditional yogurts are found, but their probiotic role was not studied. Incorporation of probiotic microorganisms (isolated from indigenous yoghurts) in market yogurts can positively enhance health status of larger segment of communities of Bangladesh.

## MATERIALS AND METHOD

### Sample and Sampling site

The samples for the study were collected from different places of Chittagong and Jessore region in Bangladesh. The experiment was conducted at the Industrial Microbiology Research Division, Bangladesh Council of Scientific and Industrial Research (BCSIR), Chittagong during the period August, 2012 to March, 2013. All the samples were collected in sterile vials. Subsequently, transferred in to sterile screw capped test tubes containing MRS broth media. At each time of collection, precaution was taken to prevent or avoid cross-contamination of samples. After collection of the samples, they were transported to the Laboratory as soon as possible in an insulated foam box with ice to maintain a temperature ranging from 4°C to 6°C. Microbiological examinations were done immediately as they arrived at the laboratory.

### Isolation and Identification of Bacteria

The collected samples were inoculated into MRS broth in conical flask and incubated at 37°C for overnight<sup>5</sup>. After overnight incubation at 37°C, opaque white colored cultures were chosen for the growth of *Lactobacillus* and *Lactococcus* sp. Culture from MRS broth (Oxoid) was then inoculated on MRS agar plates. After overnight incubation at 37°C, pure white colored colonies were tentatively chosen to be *Lactobacillus* and *Lactococcus* by observing their colony

morphology, physiological, sugar fermentation and as well as some biochemical characteristics. All the five selected strains were tested for their morphological, cultural and biochemical characters: the characters were compared with the standard description of Bergey's Manual of Systematic Bacteriology<sup>8</sup>.

### Enumeration of Bacterial Load

After incubation, the deManRogosa and Sharp Agar plates (Oxoid) were placed on a colony counter (Stuart scientific, UK) and the colonies were counted. The number of colonies or viable aerobic bacterial count per ml was calculated by multiplying the average number of colonies per plate by the reciprocal of the dilution (3). The calculated results were expressed as colony forming units (cfu) per ml of sample. McIntosh and Filde's anaerobic jar is an instrument was used in the production of an anaerobic environment.

### The Fermentation Test (SAB 1957)

To determine the products of sugar fermentation, a carbohydrate fermentation broth was prepared at pH 7.4. This nutrient broth contains 0.5% - 1.0% of the Carbohydrates ingredients to be tested (e.g. Lactose, Glucose, Sucrose, Mannose, Arabinose, Galactose, Starch and Mannitol), and the pH indicator phenol red. The nutrient broth, which is a light red color, supports the growth of most organisms whether they are able to ferment the sugar or not. The test organism is inoculated into a broth containing the test sugar and incubated at  $37 \pm 2^\circ\text{C}$  for 48 to 96 hours. A bright yellow color indicates the production of enough acid products from fermentation of the sugar to drop the pH to 6.9 or less. Production of gas is determined with a Durham tube, a small inverted vial filled with the carbohydrate fermentation broth.

### Growth Response at Different Concentration of NaCl

Nutrient broth containing different concentration of sodium chloride (0%, 3%, 4%, and 6%) was inoculated and incubated at  $37 \pm 1^\circ\text{C}$  for 48 to 72 hours. The growth of *Lactobacillus* species at different concentration of NaCl was then compared with the control.

### Preparation of Pathogenic Bacterial Suspension

One loop full 24-48 hours old pathogenic bacteria culture was taken in sterilized screw cap tube containing 2ml of sterilized saline water. The

bacterial culture was then mixed with the water thoroughly. A total of 11 pathogenic microorganisms were used in this method namely *Bacillus subtilis*, *Staphylococcus aureus*, *Streptococcus Group-B*, *Bacillus aureus*, *Escherichia coli*, *Salmonella paratyphi*, *Pseudomonas aeruginosa*, *Serratia sp.*, *Klebsiella Pneumonia*, and *Neisseria meningitidis*

#### **Activity against pathogenic microorganisms by agar well diffusion method**

In perform growth inhibition by agar well diffusion method, Mueller Hinton plates were heavily seeded ( $2.7 \times 10^3$  cells per ml) uniformly with the test organisms. Then a hole was made in media by gel cutter in sterile condition. Then one drop of melted agar was poured into hole and allowed to solidify to make a base layer. After that specific amount of culture filtrate (0.1 ml) was poured into the hole. Then plates were kept at low temperature (4°C) for 2-4 hours to allow maximum diffusion<sup>2</sup>. During this time the test materials were dissolved and diffused out of the media. The diffusion occurs according to the physical law that controls the diffusion of molecules through agar gel. As a result there is a gradual change of test materials concentration in the media surrounding the discs. The plates were then incubated at 37°C for 24 hours to allow maximum growth of the organisms at inverted position. If the test materials have any antibacterial activity, it will inhibit the growth of microorganisms giving a clear distinct zone called "zone of inhibition". The antibacterial activity of the test agent was determined by measuring the zone of inhibition expressed in mm in diameter. The experiment was carried out more than once and mean of reading is required. The selected pathogens used in the experiments were *Escherichia coli* ATCC 25922, *Streptococcus Group-B* ATCC 12401, *Staphylococcus aureus* ATCC 10832, *Haemophilus influenzae* ATCC 49766, *Klebsiella pneumoniae* ATCC 13883, *Salmonella paratyphi* ATCC 9150, *Bacillus subtilis* ATCC 21332, and *Pseudomonas aeruginosa* ATCC 15442.

#### **Determination of optimum pH and temperature for best antimicrobial activity**

The test was performed in MRS broth and the MRS broth media was adjusted for four different pH (i.e. pH 3, 3.5, 4, and 4.5) using 0.1N Acetic acid and 0.1N NaOH. Aliquots of 10 ml medium from of each pH were distributed in

separate test tube and autoclaved. After sterilization, 4 sets of tubes were inoculated with culture suspension of 5 isolates and incubated at 27°C, 37°C and 45°C temperature for 24-48 hours. After incubation the turbidity was measured with spectrophotometer at 560 nm and the culture broth were filtered with the help of Whatman filter paper (Whatman International Ltd. Maidstone, England). After filtration the final pH of the culture filtrate was recorded by pH meter (pH Hanna Instrument Ltd. & 3310, pH meter Jenway, UK) and selected for antimicrobial activity against the respective pathogenic bacteria by agar well diffusion method.

#### **Determination of Bacteriocins Production Capability of Isolates**

One ml of frozen LAB isolate was cultured overnight in 20 ml MRS broth. Then 1 ml culture was sub-cultured overnight in 20 ml MRS broth. Cells were removed by centrifuging at 9000 rpm for 15 minutes. The supernatant was filtered through a sterile Whatman No. 1 filter paper (in original paper it is 0.22 µm syringe filter) and 100 µl of the unadjusted aliquot of cell free supernatant (CFS) was added to the first well. The remaining CFS was adjusted to pH 6.0 with 1 M/1N NaOH in order to rule out possible inhibition effects due to organic acids. 100 µL of the pH adjusted CFS was filtered and added to the second well. The neutralized CFS was then treated with 1 mg/ml of catalase (Merck KGa A, Germany) at 25°C for 30 min to eliminate the possible inhibitory action of H<sub>2</sub>O<sub>2</sub> and filtered. Then 100 µl catalase CFS was placed in the third well. If inhibitions zones were found in the third well, the isolates were considered to be able to produce bacteriocin or bacteriocins like substance (BLS)<sup>17</sup>.

#### **Effect of Proteolytic Enzymes on the Antimicrobial Activity of Crude Bacteriocins (Papain and Trypsin)**

Five ml aliquot of bacteriocins was taken in test tubes and treated with papain (1 mg/ml) at pH 7. The test tubes with and without the enzyme (control) were incubated at 37°C for 2 hour and heated at 100°C for 3 minutes to denature the enzyme. Both the control and samples were assayed for antimicrobial activity by using well diffusion method (50 µL of sample in each well).

#### **Assay of Antimicrobial activity for Bacteriocin**

For testing Bacteriocin activity, after growing the selected isolates (i.e. LbD and LbP) at

optimum conditions, the culture filtrates were centrifuged at 5000 rpm for 15 minutes. The culture supernatant was then neutralized by 0.1N NaOH and the pH of the supernatant was adjusted at 7.0. Then antimicrobial activity was assayed by agar well diffusion method<sup>14</sup>.

#### Bile salt tolerance of the isolates

MRS broth was prepared with varying concentration (i.e.; 1.0%, 2.0% and 3.0%) of bile salt. Then the medium was dispensed at 10 ml per tube. Inoculation was done with the selected isolates (i.e. LcL, LcP, LcR, LbD and LbP). After inoculation with equal amount of inoculums the tubes were then incubated at 37±1°C for 24-48 hours. After incubation 0.1 ml of culture from each concentration was used to grow in agar medium by pour plate method. The plates were then incubated at 37°C for 24 hours and observed for comparative growth in these plates (4).

#### Assay of antibiotic sensitivity pattern

To assess the antibiotic sensitivity pattern disk diffusion method was followed. In this method Mueller Hinton plates were prepared and swabbed with suspension of selected isolates with the help of sterile cotton bud. After swabbing the antibiotic disks (Azithromycin, Ceftriaxone,

Amoxicillin, Kanamycin, Cefixime, Cephalexin, Streptomycin, MeropenemCefaclor, Amoxyclav, Piperacillin, Ciprofloxacin, Gentamicin, Chloramphenicol, Erythromycin, Ampicillin, Amikacin, Aztreonam, Tobramycin, Ceftazidime and Nitrofurantoin) were placed on the surface of the plate at equidistance. The plates were then kept at 4°C for 1-2 hours for proper diffusion of antibiotics. The plates were then incubated for 18-24 hours at 37°C. The zone of inhibition was observed for antibiotic sensitivity or resistance and zone diameter was measured<sup>1</sup>.

## RESULTS

#### Sugar Fermentation

Acid and Gas production by Fermentation test are given below:

#### Determination of optimum pH for best antimicrobial activity

For evaluation of the effects of pH on growth and the production of antimicrobial substances, the isolates were grown in broth medium with different pH. The growth of the organisms after incubation was measured spectrophotometrically at 560 nm and final pH of

**Table 1.** Morphological, physiological and biochemical test for identification of isolated *Lactobacilli sp*

Sample Code	Gram Staining	M. Test	T.S.I Test			I Test	MR Test	VP Test	Gel Test	O Test	Cat Test	U Test	Cit test
			Gl	Ac	H <sub>2</sub> S								
LbY-C1	+	-	+	+	+	-	+	-	+	-	-	-	+
LbY-C2	+	-	+	+	+	-	+	-	+	-	-	-	+
LbY-C3	+	-	+	+	+	-	+	-	+	-	-	-	+
LbY-C4	+	-	+	+	+	-	+	-	+	-	-	-	+
LbY-C7	+	-	+	+	-	-	+	-	+	-	-	-	+
LbY-C9	+	-	+	+	-	-	+	-	+	-	-	-	+
LbY-C11	+	-	+	+	+	-	+	-	+	-	-	-	+
LbY-C13	+	-	+	+	+	-	+	-	+	-	-	-	+
LcY-C14	+	-	+	+	+	-	+	-	+	-	-	-	+
LbY-15	+	-	+	+	+	-	+	-	+	-	-	-	+
LbY-16	+	-	+	+	+	-	+	-	+	-	-	-	+
LbY-17	+	-	+	+	-	-	+	-	+	-	-	-	+
LbY-L <sub>1</sub>	+	-	+	+	+	-	+	-	+	-	-	-	+
LbY-L <sub>9</sub>	+	-	+	+	+	-	+	-	+	-	-	-	+
LbY-J1	+	-	+	+	+	-	+	-	-	+	-	+	-
LbY-J2	+	-	+	+	+	-	+	-	-	+	-	+	-
LbY-J3	+	-	+	+	+	-	+	-	+	-	-	-	+

Note: T.S.I=Triple Sugar Iron, I= Indole, O= Oxidase, Cat= Catalase, U=Urease, Cit= Citrate, Gl=Glucose, Ac=Acidic, Gel=Gelatinase, + =Positive, - = Negative

the culture filtrate was determined. The antimicrobial activity of the culture filtrate was also assayed by agar well diffusion method.

**Determination of optimum Temperature for best antimicrobial activity**

To determine the effects of temperature

**Table 2.** Physiological and Biochemical test for identification of isolated *Lactobacilli sp*

Sample code	Growth response at different pH, Temperature & NaCl concentration (%) for <i>Lactobacilli</i> and <i>Lactococcus sp.</i> are given below:									
	Growth at different Temperature			Growth at different pH				Growth at different NaCl conc. (%)		
	27°C	37°C	45°C	3	3.5	4	4.5	3	4	6
LbY-C1	++	++	+	+	+	++	+++	++	++	+
LbY-C2	++	++	+	+	+	++	+++	++	++	+
LbY-C3	++	++	+	+	+	++	+++	++	++	+
LbY-C4	++	++	+	+	+	++	+++	++	++	+
LbY-C7	++	+	-	+	+	++	++	+	-	-
LbY-C9	++	+	-	+	+	++	++	+	+	+
LbY-C11	++	++	+	+	+	++	+++	++	++	+
LbY-C13	++	++	+	+	+	++	+++	++	++	+
LbY-C14	++	++	+	+	+	++	+++	++	++	+
LbY-15	++	++	+	+	+	++	+++	++	++	+
LbY-16	++	++	+	+	+	++	+++	++	++	+
LbY-17		++	++	-	+	+	++	++	+	-
LbY-L <sub>1</sub>	++	++	+	+	+	++	+++	++	++	+
LbY-L <sub>9</sub>	++	++	+	+	+	++	+++	++	++	+
LbY-J1	+++	+++	++	-	+	+	++	+++	++	+
LbY-J2	+++	+++	++	-	+	+	++	+++	+++	+
LbY-J3	+++	+++	+	-	+	+	++	+++	+++	+

Note: Good=+, Moderate= ++, Excellent= +++

**Table 3.** Physiological and Biochemical test for identification of isolated *Lactobacilli sp*

Sample code	Glucose	Sucrose	Lactose	Mannose	Arabinose	Galactose	Starch	Mannitol
LbY-C1	+	+	+	+	-	+	+	+
LbY-C2	+	+	+	+	-	+	+	+
LbY-C3	+	+	+	+	-	+	+	+
LbY-C4	+	+	+	+	-	+	+	+
LbY-C7	+	+	-	+	-	+	+	-
LbY-C9	+	+	+	+	+	+	+	+
LbY-C11	+	+	+	+	-	+	+	+
LbY-C13	+	+	+	+	-	+	+	+
LbY-C14	+	+	+	+	-	+	+	+
LbY-C15	+	+	+	+	-	+	+	+
LbY-C16	+	+	+	+	-	+	+	+
LbY-C17	+	+	-	+	-	+	+	-
LbY-CL <sub>1</sub>	+	+	+	+	-	+	+	+
LbY-CL <sub>9</sub>	+	+	+	+	-	+	+	+
LbY-J1	+	+	-	+	-	-	-	-
LbY-J2	+	+	-	+	-	-	-	-
LbY-J3	+	+	-	+	+	+	-	+

Note: Positive = +, Negative = -

on growth and the production of antimicrobial substances, the isolates were grown in broth medium with different temperature. The growth of the organisms after incubation was measured spectrophotometrically at 560 nm and final temperature of the culture filtrate was determined. The antimicrobial activity of the culture filtrate was also assayed by agar well diffusion method.

#### Determination of Bacteriocin Production

Zone of inhibition produced by all five isolates, indicated that all isolates are able to produce bacteriocin in our experiment. Zone of inhibition was found illustrated in the (Figure 1)

#### Effect of proteolytic enzymes on the antimicrobial activity of crude bacteriocin (Papain and Trypsin)

After treating Bacteriocin with Proteolytic enzymes papain, we have found that there were no

zone of inhibition indicated that the zone of inhibition occurred by bacteriocin. The results are shown in the below (figure 2)

#### Assay of antibiotic susceptibility

Antibiotic sensitivity pattern of the selected isolates were also determined to observe the inhibitory effect of any antibiotics against them. Combination of this type of antibiotic with probiotic organism may destroy the therapeutic activity of the probiotic. The Antibiogram obtained is shown in table-7.

From the Graphs, Isolate LbD showed 80% resistance, 20% sensitive and no intermediate found among 21 antibiotic disks. Isolate LbP showed 20% resistance, 71% sensitive and 9% intermediate among 21 antibiotic disks.

**Table 4.** Growth and final pH after incubation of the isolates at different pH


Isolates	Parameters	Initial medium pH			
		PH 3.0	PH 3.5	PH 4.0	PH 4.5
<i>Lactobacillus delbrueckii</i> (LbD)	Growth	0.327	0.421	0.489	0.89
	Final pH	3.28	3.60	4.30	4.69
<i>Lactobacillus plantarum</i> (LbP)	Growth	0.320	0.396	0.770	0.96
	Final pH	3.28	3.80	4.50	4.8

**Table 5.** Growth and final pH after incubation of the isolates at different temperature

Isolates	Parameters	Incubation Temperature		
		27°C	37°C	45°C
<i>Lactobacillus delbrueckii</i> (LbD)	Growth	0.195	0.210	0.177
	Final pH	1.09	1.08	1.13
<i>Lactobacillus plantarum</i> (LbP)	Growth	0.206	0.215	0.183
	Final pH	1.04	1.03	1.05

**Table 6.** Assay of Bacteriocin activity produced by the selected isolates given below


Isolates	Pathogenic bacteria	Zone of inhibition
<i>Lactobacillus delbrueckii</i> (LbD)	<i>Escheriachia coli</i>	22
	<i>Streptococcus Group-B</i>	14
	<i>Staphylococcus aureus</i>	18
	<i>Haemophilus influenza</i>	16
	<i>Klebsiella Pneumonia</i>	12
<i>Lactobacillus plantarum</i> (LbP)	<i>Escheriachia coli</i>	24
	<i>Streptococcus Group-B</i>	20
	<i>Staphylococcus aureus</i>	22
	<i>Haemophilus influenza</i>	20
	<i>Klebsiella Pneumonia</i>	12




**Fig. 1.** Determination of Bacteriocin Production Capability of the Lab Isolates




**Fig. 2.** Effect of Proteolytic Enzymes on the Antimicrobial Activity of Crude Bacteriocin (Papain and Trypsin)




**Fig. 3.** Percentage of Antibiotic Resistance, 21 for *Lactobacilli sp.*



**Fig. 4.** Zone of inhibition produced produced by isolate LbD and LbP against 05 pathogenic organisms



5




6

**Fig. 5-6.** Zone of inhibition against (5) *E. coli* and (6) *Streptococcus Group B*



7



8

**Fig. 7-8.** Antibiotic sensitivity and resistant of (7) *Lactobacillus delbrueckii* (8) *Lactobacillus plantarum*

**Assay of antimicrobial activity for bacteriocin**

Bacteriocin an antimicrobial substance produced by most lactic acid bacteria. The activity of Bacteriocin produced by the selected isolates was also determined and the effectively of the substance was measured by the zone of inhibition (mm) which given following table-6.

In the Graph, LbD and LbP produced zone of inhibition (mm) respectively 22 and 24 mm against pathogenic bacteria *E. coli.*, and 14, 20 mm against pathogenic bacteria *Streptococcus Group-B.*, and 18, 22 mm against pathogenic bacteria *Staphylococcus aureus.*, and 16, 20 mm against pathogenic bacteria *Haemophilus influenza.*, and 13, 12 mm against pathogenic bacteria *Klebsiella pneumonia.*

**Bile salt tolerance**

As probiotic organism has to tolerate different bile salt concentration in the intestinal tract or stomach for their survival, the isolates were assayed for their sensitivity to different bile salt


concentration (i.e. 1%, 2% and 3%). The activities they showed were tabulated-12 in the following. In the graph showed that when bile salt concentration is increased, the No. of colony is decreased. Every isolate tolerate up to 3% bile salt but 1% bile salt gives the best no. of colony (CFU/0.1ml) for all isolate.

**DISCUSSION**

For isolation of lactic acid bacteria, yoghurt samples were collected from different places of Chittagong & Jessore region. The identification of isolates were determined according to the Bergey's Manual of Systematic Bacteriology (2009). pH is an important factor which can dramatically affect bacterial growth. In our experimental design we have observed the growth of our isolated lactobacillus in various pH values ranges (3 to 4.5). The result shown that all the isolates grew best at pH 4.5 used range value (Table-5). Growth increases with the increasing of pH from pH-3 to pH-4.5 but growth decrease when it exceeds the neutral range of pH. It was found that *Lactobacillus delbrueckii* subsp. *Bulgaricus*, *Lactococcus lactis* subsp. *Lactis*, *Lactococcus raffinolactis*, *Enterococcus faecium*, *Pediococcus pentosaceus* can tolerate pH up to 5.0 %. The present results are closely related to their findings. It was reported that *Lactobacillus plantarum* could produce lactic acid and reduce the pH to values lower than 4.0 (15). After growth at different temperature i.e. 27°C, 37°C and 45°C, it was found that the isolates grew best at 37°C but *Lactobacillus delbrueckii* and *Lactobacillus plantarum* can grow at 45°C and show suitable for best antimicrobial activity (Table-7). The growth decreases with the increasing temperature (i.e.; 45°C). NaCl is an inhibitory substance which may inhibit growth of certain types of bacteria. The current results showed that *Lactobacillus delbrueckii* and *Lactobacillus plantarum* were able to tolerate 3%, 4%, & 6% NaCl and good growth was observed at 4% NaCl (Table-8). It was found that 56% of Lactococci tested were able to grow on 6.5% NaCl <sup>14</sup>. It was reported that *Lactococcus lactis* was the most tolerant to high NaCl concentration compared to the other isolates. The table shows the number of colonies of the isolates on agar medium after incubating them in



**Fig. 9.** Antibiotic sensitivity and resistant of *Lactobacillus plantarum* (LbP)



In the graph showed that when bile salt concentration is increased, the No. of colony is decreased. Every isolate tolerate up to 3% bile salt but 1% bile salt gives the best no. of colony (CFU/0.1ml) for all isolate.

**Fig. 10.** Number of colony count for isolates against different bile salt concentration (%)

MRS broth medium containing varying concentration of bile salt i.e. 1.0 %, 2.0 %, 3.0%. In the Table-8, it is seen that at lower concentration the colony numbers were very high but gradually

**Table 7.** Antibiotic Sensitivity pattern of the selected isolates given below

Antibiotics disk (µg)	<i>Lactobacillus delbrueckii</i> LbD (mm)	<i>Lactobacillus plantarum</i> LbP(mm)
GEN-30	R	30 S
CFM-5	R	10 R
CP-30	R	25 S
E-15	R	24 S
CI-30	-	-
AZM-30	R	26 S
K-30	R	21 S
AMX-30	-	-
C-30	28 S	20 S
MRP-10	R	37 S
CIP-5	R	40 S
AML-10	R	18 S
PI-100	28 S	30 S
CF-30	R	13 I
AC-30	30 S	40 S
S-10	21 S	22 S
A-25	R	10 R
AK-30	R	28 S
ATM-30	R	R
TOB-10	R	20 S
VA-30	R	20 S
CAZ-30	R	R
F-300	R	14 I

Note: - = Not done, R= Resistent, I = Intermediate, S = Sensitive, GEN = Gentamicin, AZM = Azithromycin, E= Erythromycin, CI = Ceftriaxome, K= Kanamycin, CP= Cephalexin, CFM= Cefixime, S = Streptomycin, MRP = Meropenem, CF = Cefaclor, AML = Amoxicillin, C = Chloramphenicol, AC= Amoxyclav, CIP = Ciprofloxacin, PI = Piperacillin, A = Ampicillin, AK = Amikacin, ATM = Aztreonam, TOB = Tobramycin, VA = Vanomycin, CAZ = Ceftazidime, F = Nitrofurantoin.

the numbers decreased with the increasing of bile salt concentration. The zone of inhibition in the Table-6 indicates that the isolates secrete Bacteriocin and it has bactericidal activities. The table-6 shows that Zone of inhibition (mm) produced by *L. delbrueckii*, *L. plantarum*, against pathogenic bacteria *E. coli*, *Streptococcus Group-B*, *Staphylococcus aureus*, *Haemophilus influenza*, *Klebsiella pneumonia*, *Salmonella paratyphi*, *Bacillus subtilis*, and *Pseudomonas aeruginosa* were satisfactory. Inhibition by isolates and further treated with papain indicate that our isolates inhibited the growth of pathogenic microorganisms by producing Bacteriocins like substances. Our study showed similarity with other research such results have shown *L. plantarum* higher antimicrobial activities against *E. coli* and *Staph. aureus* than the others. The zone of inhibition by *L. plantarum* against *E. coli* were 12.33 mm while zone of inhibition against *Staph. aureus* were 14.33 mm but it showed *L. plantarum* was able to inhibit the growth of the Gram negative tested bacteria with an average inhibition zone of 18 and 26 mm in diameter against *Salmonella* sp. and *E. coli* strains respectively<sup>17</sup>. It was reported *Lactococcus lactis ssp. lactis* antagonistic behavior was demonstrated by the production of 14 mm against *Salmonella* sp. and 20 mm inhibition zones against *E. coli* strains which supports our findings<sup>16</sup>.

Lactic acid bacteria (LAB) from fermented products may act as a reservoir of antimicrobial-resistance genes<sup>7</sup>. *Lactobacillus delbrueckii* were found sensitive to Chloramphenicol, Piperacillin, Amoxyclav and Streptomycin but resistant to remaining 17 antibiotics. Our results were similar to a study<sup>3</sup> *Lactobacillus plantarum* is Sensitive to 15 antibiotics used against them and Intermediate to Cefaclor, Nitrofurantoin but resistant to Cefixime, Ampicillin, Aztreonam and Ceftazidime. From the seven isolates of *Lactobacillus plantarum* 3

**Table 8.** Bile salt tolerance (at different concentrations) of the selected isolates given below

Isolates	No. of colony grown on the plates (CFU/0.1 ml)		
	1.0%	2.0%	3.0%
<i>Lactobacillus delbrueckii</i> (LbD)	360	220	180
<i>Lactobacillus plantarum</i> (LbP)	510	450	350

isolates were resistant to Ampicillin and one resistant to tetracycline. *Lactobacilli* isolated from milk and milk products were resistant to erythromycin, Ampicillin and tetracycline and were sensitive in 100 % to Gentamicin which is in agreement with our findings. Such resistance to a wide spectrum of antibiotic therapy may be helpful in faster recovery of the patients due to rapid establishment of desirable microbial flora. This study supports the use of selected probiotic agents for the prevention of antimicrobial-associated diarrhea. Resistance of the probiotic strains to some antibiotics could be used both preventive and therapeutic purpose in controlling intestinal infections<sup>4</sup>.

In conclusion, the experimental results showed that isolated two selected species were able to tolerate inhibitory substance bile salt at 1-3% where best at 1% and 3-6% NaCl. The suitable temperature for their growth is showed 37°C and pH-4.5, but *Lactobacillus delbrueckii* and *Lactobacillus plantarum* grew at temperature 45°C. Growth inhibition against selected pathogens (*Escherichia coli* ATCC 25922, *Streptococcus Group-B* ATCC 12401, *Staphylococcus aureus* ATCC 10832, *Haemophilus influenzae* ATCC 49766, *Klebsiella pneumoniae* ATCC 13883, *Salmonella paratyphi* ATCC 9150, *Bacillus subtilis* ATCC 21332, and *Pseudomonas aeruginosa* ATCC 15442 and *Neisseria meningitidis* ATCC 35561) and resistance against various antibiotics suggest that our identified *Lactobacilli* (*Lactobacillus delbrueckii* and *Lactobacillus plantarum*). On the other hand *Lactobacillus delbrueckii* was resistant to 17 antibiotics used against them but *Lactobacillus plantarum* were found resistant only against 04 antibiotics (Cefixime, Ampicillin, Aztreonam and Ceftazidime). Finally, it can be said that the selected *lactobacillus delbrueckii* found in our study showed potentially to be used as a probiotics in near future. In spite of the problems with dosage and viability of probiotic strains, lack of industry standardization and potential safety issues, there is obviously considerable potential for the benefits of probiotics over a wide range of clinical conditions. Further study on these isolates will help to detect the genes responsible for therapeutic activities and this type of research will help to design more probiotic agents to control numerous

diseases and at the same time it will ensure the safe and healthy human civilization.

#### ACKNOWLEDGMENTS

We thank Bangladesh Council of Scientific and Industrial Research (BCSIR) for their logistic support and collaboration.

#### Conflict of Interest

We hereby declare that we have no conflict of interest regarding this paper

#### REFERENCES

1. Bacha, K., Mehari, T. & Ashenafi, M. In-vitro probiotic potential of lactic acid bacteria isolated from 'Wakalim', a traditional Ethiopian fermented beef sausage. *Ethiop J Health Sci*, 2009; **19**: 21-29.
2. Bauer, A., Kirby, W., Sherris, J. C., Turck & Turck, M., Antibiotic susceptibility testing by a standardized single disk method. *American journal of clinical pathology*, 1966; **45**: 493.
3. Belletti, N., Gatti, M., Bottari, B., Neviani, E., Tabanelli, G. & Gardini, F., Antibiotic resistance of lactobacilli isolated from two Italian hard cheeses. *Journal of Food Protection*®, 2009; **72**: 2162-2169.
4. Bhutada, S. & Tambekar, D., An evaluation of probiotic potential of *Lactobacillus* sp. from milk of domestic animals and commercial available probiotic preparations in prevention of enteric bacterial infections. *Recent Research in Science and Technology*, 2010; **2**.
5. Collins, C. H., Microbiological methods. *Microbiological methods*, 1964.
6. Diop, M., Dubois Dauphin, R., Dortu, C., Destain, J., Tine, E. & Thonart, P., In vitro detection and characterization of bacteriocin-like inhibitory activity of lactic acid bacteria (LAB) isolated from Senegalese local food products. *African Journal of Microbiology Research* [= *AJMR*], 2008; **2**.
7. Flórez, A. B., Delgado, S. & Mayo, B., Antimicrobial susceptibility of lactic acid bacteria isolated from a cheese environment. *Canadian journal of microbiology*, 2005; **51**: 51-58.
8. Garrity, G. M., Bell, J. A. & Lilburn, T. G., Taxonomic outline of the prokaryotes. *Bergey's manual of systematic bacteriology*. Springer, New York, Berlin, Heidelberg, 2004.
9. Harun-ur-rashid, M., Togo, K., Ueda, M. & Miyamoto, T., Probiotic characteristics of lactic

- acid bacteria isolated from traditional fermented milk 'Dahi' in Bangladesh. *Pakistan J Nutr*, 2007; **6**: 647-652.
10. Heczko, P. B., Strus, M. & Kochan, P., Critical evaluation of probiotic activity of lactic acid bacteria and their effects. *J Physiol Pharmacol*, 2006; **57**(9): 5-12.
  11. Kumar, M., Kumar, A., Nagpal, R., Mohania, D., Behare, P., Verma, V., Kumar, P., Poddar, D., Aggarwal, P. K., Henry, C. J., Jain, S. & Yadav, H., Cancer-preventing attributes of probiotics: an update. *Int J Food Sci Nutr*, 2010; **61**: 473-96.
  12. Liu, C. F., Tseng, K. C., Chiang, S. S., Lee, B. H., Hsu, W. H. & Pan, T. M., Immunomodulatory and antioxidant potential of *Lactobacillus* exopolysaccharides. *Journal of the Science of Food and Agriculture*, 2011; **91**: 2284-2291.
  13. Murray, B. E., The life and times of the Enterococcus. *Clinical microbiology reviews*, 1990; **3**: 46-65.
  14. Parente, E., Brienza, C., Moles, M. & Ricciardi, A., A comparison of methods for the measurement of bacteriocin activity. *Journal of microbiological methods*, 1995; **22**: 95-108.
  15. Rao, M. S., Pintado, J., Stevens, W. F. & Guyot, J. P., Kinetic growth parameters of different amyolytic and non-amyolytic *Lactobacillus* strains under various salt and pH conditions. *Bioresour Technol*, 2004; **94**: 331-7.
  16. Yateem, A., Balba, M., Al-surrayai, T., Al-mutairi, B. & Al-daher, R., Isolation of Lactic Acid Bacteria with Probiotic Potential from Camel Milk. *International Journal of Dairy Science*, 2008; **3**.
  17. Yang, E., Fan, L., Jiang, Y., Doucette, C. & Fillmore, S., Antimicrobial activity of bacteriocin-producing lactic acid bacteria isolated from cheeses and yogurts. *AMB Express*, 2012; **2**: 1-12.
  18. Zourari, A., Accolas, J. & Desmazeaud, M., Metabolism and biochemical characteristics of yogurt bacteria. A review. *Le lait*, 1992; **72**: 1-34.