

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/333140926>

Isolated Endophytic Fungi from the Plant *Curcuma longa* and Their Potential Bioactivity—A Review

Article in *Pharmacology & Pharmacy* · January 2019

DOI: 10.4236/pp.2019.105021

CITATIONS

5

READS

404

5 authors, including:



Kazi Jannatul Ferdous

North South University

2 PUBLICATIONS 6 CITATIONS

[SEE PROFILE](#)



Farhana Afroz

Bangladesh Council of Scientific and Industrial Research

45 PUBLICATIONS 156 CITATIONS

[SEE PROFILE](#)



Md Abdul Mazid

University of Dhaka

82 PUBLICATIONS 591 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Emerging and re-emerging infectious diseases in Bangladesh [View project](#)



1 Chemical exploration and value addition in bioactive molecules isolated from medicinal plants and their associated endophytic fungi. [View project](#)

Isolated Endophytic Fungi from the Plant *Curcuma longa* and Their Potential Bioactivity—A Review

Kazi Jannatul Ferdous^{1,2}, Farhana Afroz², Md. Rakibul Islam¹, Md. Abdul Mazid³,
Md. Hossain Sohrab^{2*}

¹Department of Biochemistry and Molecular Biology, University of Dhaka, Dhaka, Bangladesh

²Pharmaceutical Sciences Research Division, Bangladesh Council of Scientific and Industrial Research, Dhaka, Bangladesh

³Department of Pharmaceutical Chemistry, University of Dhaka, Dhaka, Bangladesh

Email: *psrd_bcsir@yahoo.com

How to cite this paper: Ferdous, K.J., Afroz, F., Islam, M.R., Mazid, M.A. and Sohrab, M.H. (2019) Isolated Endophytic Fungi from the Plant *Curcuma longa* and Their Potential Bioactivity—A Review. *Pharmacology & Pharmacy*, 10, 244-270. <https://doi.org/10.4236/pp.2019.105021>

Received: April 9, 2019

Accepted: May 14, 2019

Published: May 17, 2019

Copyright © 2019 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Endophytes (both fungi and bacteria) represent a treasure house for bioactive compounds such as anticancer, immunomodulatory, antioxidant, antiparasitic, insecticidal etc. for use in the pharmaceutical and agrochemical industries. Endophytic population is greatly influenced by climatic conditions and location where the host plant grows. *Curcuma longa* L. (Turmeric) is a medicinal plant belonging to the Zingiberaceae family that is usually used as spice and preservative. Traditionally it is used to treat various conditions and symptoms such as digestive disorders, arthritis, cardiovascular conditions, cancer, bacterial infections etc. In recent times, there have been different discoveries on endophytic fungi and their biological activity from *C. longa*. But there is no comprehensive review on endophytic fungi of *C. longa*. The aim of this review is to analyze the published report based on the endophytic fungi to provide for the first time an updated information about their isolation from different tissues, their biological activities including the phytochemistry of *C. longa*. This will lead the way to forecast the missing link for future research work.

Keywords

Curcuma longa, Endophytes, Compounds, Bioactivity

1. Introduction

Curcuma longa L. (Turmeric) is a rhizomatous herbaceous perennial plant of the

ginger family, Zingiberaceae [1]. It is a naturally grown plant in the Indian sub-continent and Southeast Asia. Turmeric has been used in Asia for thousands of years and is a major part of Ayurveda, Siddha medicine, Unani, and traditional Chinese medicine [2]. A lot of scientific works on pharmacological properties of turmeric have been reported by many scientists and researchers such as anti-inflammatory [3] [4], antiallergic [5], antiobesity [6], lipid-lowering [7], anti-proliferative [8], anti-microbial [9] [10], anti-oxidant [11] [12], anti-diabetic [13], hypoglycemic [14], immunomodulating [15], insecticidal [16], anti-fungal [17], neuroprotective [18], anti-mutagenic [19], hypolipidemic [20] etc.

Endophyte, the most important integral part of plant tissues, is a progressively significant area of research in many fields because of their chemical miscellany and their ability to produce many unique secondary metabolites that can be utilized for fuel, medicine, restoration and agriculture. Endophytes are isolated from medicinal plants possess strong fungicidal, bactericidal and cytotoxic metabolites [21]. Previous studies indicate that there have been few investigations done on endophytic microbes of turmeric plant (*Curcuma longa*). Thus, exploring endophytic microbes from *Curcuma longa* increases the probability to discover novel compound which is extensive in resistance tissues. There is no review on the endophytic fungi from turmeric plants. Therefore, it is necessary to review the endophytic fungi from *Curcuma longa*. Hence, this review discusses the current research progresses on endophytic fungi from the turmeric plant, including types of endophytic fungi identified, described bioactive compounds, and biological activities of endophytes to discover the current scope of knowledge of endophytes.

2. Isolated Compounds from *Curcuma longa*

A total of 235 compounds with various activities were isolated and identified from the flowers, leaves, roots and rhizomes part of *Curcuma longa* which is the most phytochemically investigated plant in the world. These compounds are primarily phenolic and terpenoid compounds including diarylheptanoids (including commonly known as curcuminoids), diarylpentanoids, monoterpenes, sesquiterpenes, diterpenes, triterpenes, sterols, fatty acids and also some alkaloidal compounds. A list of isolated compounds from different groups of *C. longa* is presented in Table 1. Curcumin is the key phytochemical compound

Table 1. Isolated compounds from *C. longa*.

Sl.No	Compounds	Compound type	References
1	curcumin (curcumin I), demethoxycurcumin (curcumin II), 1-(4-hydroxy-3-methoxyphenyl)-7-(3,4-dihydroxyphenyl)-1,6-heptadiene-3,5-dione, 1-(4-hydroxyphenyl)-7-(3,4-dihydroxyphenyl)-1,6-heptadiene-3, 5-dione,	Diarylheptanoid	[24] [28]-[35]

Continued

	bisdemethoxycurcumin (curcumin III), tetrahydroxycurcumin, 5-hydroxyl-1-(4-hydroxy-3-methoxyphenyl)-7-(4-hydroxyphenyl)-4,6-heptadiene-3-one, 5-hydroxyl-1,7-bis(4-hydroxy-3-methoxyphenyl)-4,6-heptadiene-3-one, 1,7-bis(4-hydroxyphenyl)-1-heptene-3,5-dione, 5-hydroxyl-7-(4-hydroxy-3-methoxyphenyl)-1-(4-hydroxyphenyl)-4,6-heptadiene-3-one, 3-hydroxy-1,7-bis-(4-hydroxyphenyl)-6-heptene-1,5-dione, 1,5-dihydroxy-1-(4-hydroxy-3-methoxyphenyl)-7-(4-hydroxyphenyl)-4,6-heptadiene-3-one, 1,5-dihydroxy-1-(4-hydroxyphenyl)-7-(4-hydroxy-3-methoxyphenyl)-4,6-heptadiene-3-one, 1,5-dihydroxy-1,7-bis(4-hydroxy-3-methoxyphenyl)-4,6-heptadiene-3-one, 1,5-dihydroxy-1,7-bis(4-hydroxyphenyl)-4,6-heptadiene-3-one, 1,5-epoxy-3-carbonyl-1,7-bis(4-hydroxyphenyl)-4,6-heptadiene, Cyclocurcumin, 1,7-bis(4-hydroxy-3-methoxyphenyl)-1,4,6-heptatrien-3-one, 1,7-bis-(4-hydroxyphenyl)-1,4,6-heptatrien-3-one, 1,5-bis(4-hydroxyphenyl)-penta-(1E,4E)-1,4-dien-3-one, 1-(4-hydroxy-3-methoxyphenyl)-5-(4-hydroxyphenyl)-1,4-pentadiene-3-one, 1,5-bis(4-hydroxy-3-methoxyphenyl)-penta-(1E,4E)-1,4-dien-3-one		
2	4''-(4'''-hydroxyphenyl)-2''-oxo-3''-butenyl-3-(4'-hydroxyphenyl-3'-methoxy)-propenoate, 4''-(4'''-hydroxyphenyl-3-methoxy)-2''-oxo-3''-butenyl-3-(4'-hydroxyphenyl)-propenoate, calebin-A, (E)-4-(4-hydroxy-3-methoxyphenyl)but-3-en-2-one, (E)-ferulic acid, (Z)-ferulic acid	Phenylpropene	[28] [35] [36]
3	vanillic acid, vanillin	Phenolic	[35]
4	p-cymene, m-cymene, α -terpinene,	Monoterpenoid	[27] [38]-[44]

Continued

γ -terpinene,
 β -phellandrene,
p-mentha-1,4(8)-diene,
terpinen-4-ol,
4-terpinol,
Limonene,
Terpinolene,
Thymol,
Phellandrol,
Carvacrol,
(E)-carveol,
 γ -terpineol,
Menthol,
1,3,8-paramenthatriene,
p-methylacetophenone,
Piperitone,
o-cymene,
Carvone,
p-menth-8-en-2-one,
 α -thujene,
 α -terpineol,
p-cymen-8-ol,
p-meth-8-en-2-one,
piperitone epoxide,
Sylvestrene,
Menthofuran,
 β , β -dimethylstyrene,
Camphor,
Teresantalol,
Benzene,
1-methyl-4-(1-methylpropyl),
2-norpinanone,
borneol,
bornyl acetate,
(E)-chrysanthenyl acetate,
(Z)-cinerone,
(Z)-sabinol,

Continued

	2-(2,5-dihydroxy-4-methylcyclohex-3-enyl) propanoic acid, camphene, 3-carene, 2-carene, ascaridole, α -pinene, β -pinene, cineole, cis-ocimene, citronellal, geranial, neral, myrcene, R-citronellene, citronellylpentanoate, nerol, geraniol, iso-artemisia ketone, trans-ocimene, linalool, neryl acetate, geranic acid, geranyl acetate, 3-bornanone, 4,8-dimethyl-3,7-nonadien-2-ol, 3,4,5,6-tetramethyl-2,5-octadiene, 3,7-dimethyl-6-nonenal, 2,6-dimethyl-2,6-octadiene-1,8-diol, 4,5-dimethyl-2,6-octadiene		
5	ar-turmerone, α -turmerone, β -turmerone, 2-methyl-6-(4-hydroxyphenyl)-2-hepten-4-one, 2-methyl-6-(4-hydroxy-3-methylphenyl)-2-hepten-4-one, 2-methoxy-5-hydroxybisabola-3,10-diene-9-one,	Bisabolane	[25] [26] [35]-[52]

Continued

2-methyl-6-(4-formylphenyl)-2-hepten-4-one,
 5-hydroxyl-ar-turmerone,
 4-methylene-5-hydroxybisabola-2,10-diene-9-one,
 ar-curcumene,
 ar-turmerol,
 bisabola-3,10-diene-2-one,
 bisabolone,
 4, 5-dihydroxybisabola-2,10-diene,
 4-hydroxybisabola-2,10-diene-9-one,
 4-methoxy-5-hydroxy-bisabola-2,10-diene-9-one,
 bisacurone,
 bisacurone A,
 bisacurone B,
 bisacurone C,
 bisabolone-9-one,
 bisacumol,
 turmeronol A,
 turmeronol B,
 α -oxobisabolene,
 α -zingiberene,
 xanthorrhizol,
 zingerone,
 dehydrozingerone,
 (Z)- α -atlantone,
 (E)- α -atlantone,
 β -bisabolene,
 (6S,7R)-bisabolene,
 γ -bisabolene,
 γ -curcumene,
 β -curcumene,
 α -curcumene,
 β -sesquiphellandrene,
 (Z)- γ -atlantone,
 (E)- γ -atlantone,
 (6S)-2-methyl-6-[(1R,5S)-(4-methene-5-hydroxyl-2-cyclohexen)-2-hepten-4-one,
 curcuphenol,
 curlone,

Continued

	curculonone C, curculonone D, curculonone B, curculonone A, 2, 5-dihydroxybisabola-3, 10-diene, (6R)-[(1R)-1,5-dimethylhex-4-enyl]-3-methylcyclohex-2-en-1-one, β -atlantone, 2,8-epoxy-5-hydroxybisabola-3,10-diene-9-one, α -bisabolol, dihydro-ar-turmerone, dehydrocurcumene		
6	(4S,5S)-germacrone-4,5-epoxide, dehydrocurdione, germacrene D, germacrone, germacrone-13-al, β -germacene	Germacrane	[39] [40] [49]
7	1,10-dehydro-10-deoxy-9-oxozedoaronediol, curcumenol, epiprocurcumenol, isoprocurcumenol, zedoaronediol, procurcumadiol, procurcumenol	Guaiane	[35] [49]
8	naphthalene, 1,2,3,4,4a,5,6,8a-octahydro-4a,8-dimethyl-2-(1-methylethylidene), α -selinene, juniper camphor, corymbolone	Selinane	[38] [41]
9	α -santalol, α -santalene, β -santalene	Santalane	[38]
10	(E)-caryophyllene, caryophyllene oxide	Caryophyllane	[38] [43]
11	β -elemene, γ -elemene	Elemene	[38] [41]
12	acoradiene	Acorane	[38]
13	aristolene	Aristolene	[38]
14	(Z)- α -bergamotene	Bergamotane	[43]

Continued








15	curcumenone	Carabrane	[49]
16	di-epi-cedrene	Cedrane	[38]
17	himachalene	Himachalene	[38]
18	(E)-sesquisabinene hydrate	Sesquisabinane	[43]
19	bicyclo [7.2.0] undecane,10,10-dimethyl-2,6-bis(methylene), 10,10-dimethyl-2,6-bis(methylene), γ -gurjunen epoxide, 1-epi-cubenol, cubebene, 7-epi-sesquithujene, caryophyllene, 6 α -hydroxycurcumanolide A, curcumanolide A, curcumanolide B, curcumin L, α -humulene, 12-oxabicyclo [9.1.0]dodeca-3,7-diene, 1,5,5,8-tetramethyl, adoxal, 2,6,10-dodecatrien-1-ol, 3,7,11-trimethyl, (E,E)- α -farnesene, 5,9-undecadien-2-one, 6,10-dimethyl-, (Z), hxadecane-1,2-diol, nerolidal, (Z)- β -farnesene, nerolidyl propionate	Sesquiterpene	[35] [37] [38] [39] [40] [41] [43] [51] [53]
20	Phytol, (E,E,E)-3,7,11,15-tetramethylhexadeca- 1,3,6,10,14-pentaene, 2,6,11,15-tetramethyl-hexadeca-2,6,8,10,14-pentaene, 1,6,10,14-hexadecatetraen-3-ol, 3,7,11,15-tetramethyl-, (E,E)	Diterpenoid	[38] [41]
21	hopenone I, hop-17(21)-en-3 β -ol, hop-17(21)-en-3 β -yl acetate	Triterpenoid	[54]

Continued


22	β -sitosterol, stigmasterol, gitoxygenin, 20-oxopregn-16-en-12-yl acetate	Steroid	[35] [38]
23	linoleic acid, 8,11-Octadecadienoic acid, methyl ester, palmitic acid (n-hexadecanoic acid), oleic acid, stearic acid	Fatty acid	[41]
24	curcuma-J, 2-(2'-methyl-1'-propenyl)-4,6- dimethyl-7-hydroxyquinoline, 2,3,5-trimethylfuran, (1,2,3-trimethyl-cyclopent-2-enyl)-methanol, dicumyl peroxide, 1-(3-cyclopentylpropyl)-2,4-dimethyl-benzene, 1,4-dimethyl-2-(2-methylpropyl)-benzene, 2,2'-oxybis [octahydro-7,8,8-trimethyl- 4,7-methanobenzofuran, cyclohexylformate, methyleugenol, 3,3,5-trimethyl-cyclohexanol acetate, 2,4-dimethyl-8-oxabicyclo [3.2.1]oct-6-en-3-one, 2,6-dimethyl-6-(4-methyl-3-pentenyl)-2- cyclohexene-1-carboxaldehyde, bicyclo [3.3.1]nonan-9-one, 2,4-dimethyl-3-nitro-(exo), 2,2,4-trimethyl-3-(3,8,12,16-tetramethyl-heptadeca- 3,7,11,15-tetraenyl)-cyclohexanol, pyrazolo [1,5-a]pyridine, 3,3a,4,7-tetrahydro-3,3-dimethyl-, (3aS)	Miscellaneous	[38] [50] [55]

responsible for many of turmeric's eccentric properties and bright yellow colouring. Curcumin has been shown to reveal antioxidant, anti-inflammatory, antiviral, antibacterial, antifungal, and anticancer activities and thus, has a potential against various malignant diseases, diabetes, allergies, arthritis, Alzheimer's disease, and other chronic illnesses [22]. Ar-turmerone and curlone have antimutagenic activity [2] [23] whereas sodium curcuminates has anti-inflammatory activity [2]. The structures of some bioactive compounds from *C. longa* are given in **Table 2**.

Table 2. Structures of some bioactive compounds from *C. longa*.


Compound type	References
Diarylheptanoid	
 curcumin I	[28] [29]
 curcumin II	[28] [29]
 curcumin III	[28]
 tetrahydrocurcumin	[28]
 1,7-bis(4-hydroxyphenyl)-1-heptene-3,5-dione	[31]
Phenylpropene	
 4''-(4'''-hydroxyphenyl)-2''-oxo-3''-butenyl-3-(4'-hydroxyphenyl-3'-methoxy)-propenoate	[36]
 4''-(4'''-hydroxyphenyl-3-methoxy)-2''-oxo-3''-butenyl-3-(4'-hydroxyphenyl)-propenoate	[36]

Continued




[28]

calebin-A



[35]


(E)-4-(4-hydroxy-3-methoxyphenyl)but-3-en-2-one



[35]


(E)-ferulic acid

Phenolic



[35]


vanillic acid



[35]


vanillin

Monoterpenoid



[40]


thymol



[38]


menthol

Continued




piperitone

[38]




geraniol

[38]




geranic acid

[38]


Bisabolane*ar*-turmerone

[25] [47]




zingerone

[34]



curlone

[35]




2,5-dihydroxybisabola-3,10-diene

[34]


Continued

Germacrane




[49]

(4S,5S)-germacrone-4,5-epoxide




[49]

dehydrocurdione




[39]

germacrene D



[49]


germacrone



[49]


germacrone-13-al

Guaiane




[49]

epiprocurcumenol



[49]


isoprocurcumenol



[49]


zedoaronediol

Continued




[49]

procurcumadiol




[49]

procurcumenol


Selinane

[38]

naphthalene, 1,2,3,4,4a,5,6,8a-octahydro-4a,8-dimethyl-2-(1-methylethylidene)




[38]

 α -selinene


[38]

juniper camphor



[38]


corymbolone

Santalane

[38]


 α -santalol

Continued



[38]


α -santalene



[38]


β -santalene

Caryophyllane



[43]


(E)-caryophyllene



[38]


caryophyllene oxide

Elemene



[41]


β -elemene



[38]

γ -elemene


Acorane



[38]


acoradiene

Continued


Aristolene

[38]

aristolene


Bergamotane

[43]

(Z)- α -bergamotene**Carabrane**


[49]

curcumenone

Cedrane


[38]

di-epi-cedrene

Himachalene


[38]

himachalene

Sesquisabinane

[43]


(E)-sesquisabinene hydrate

Sesquiterpene

[43]


7-epi-sesquithujene

Continued




caryophyllene

[39]




adoxal

[38]




nerolidal

[37]




nerolidyl propionate

[38]

Diterpenoid


phytol

[41]




(E,E,E)-3,7,11,15-tetramethylhexadeca-1,3,6,10,14-pentaene

[38]




2,6,11,15-tetramethyl-hexadeca-2,6,8,10,14-pentaene

[38]



1,6,10,14-hexadecatetraen-3-ol,3,7,11,15-tetramethyl-,(E,E)-


[38]

Triterpenoid


hopenone I

[54]


Continued




[54]




[54]

Steroid


[35]



[35]


Fatty acid

[41]




[41]

Continued



[41]


oleic acid



[41]


stearic acid

Miscellaneous




[55]

curcuma-J




[34]

2-(2'-methyl-1'-propenyl)-4,6-dimethyl-7-hydroxyquinoline




[38]

2,3,5-trimethylfuran



[38]

methyleugenol



[38]

3,3,5-trimethyl-cyclohexanol acetate

3. Isolated Endophytic Fungi from *Curcuma longa*

Endophytic fungi *Fusarium nivale*, *Fusarium solani*, *Doratomyces stemonitis*, *Penicillium*, *Diaphorte sp.* were isolated from *Curcuma longa* [56] [57] [58]. Various studies have been identified the prospective of endophytic fungi from this plant as antioxidant, antidiabetic, anti-microbial and antitumor [62] [63]

[64] [65] [66]. Therefore, in this section, endophytic fungi from different parts of *C. longa* is reviewed and presented in **Table 3**.

3.1. Endophytic Fungi of Rhizome

From the rhizome of *C. longa* several endophytic fungi have been isolated [62]. Exactly 207 endophytic fungi were found to be isolated in one of the study that belongs to the seven genera namely, *Acremonium*, *Aspergillus*, *Exophiala*, *Fusarium*, *Penicillium*, *Phoma*, and *Stachybotrys*. Among the species *F. solani*, *F. delphinoides*, *F. verticillioides* and *A. versicolor*, *A. venenatus* bearing the genus of *Fusarium sp.* and *Aspergillus sp.* were recovered from rhizomes of *C. longa*. The rest of the genera were isolated as single species. From them, *Exophiala lecanii* the fungal endophyte was isolated for the first time from the rhizome of *C. longa* [59]. Another investigation reported 31 endophytic fungi from the rhizome of *C. longa* [60]. In most studies of endophytic fungi, the sterile mycelia fungi are reported as mycelia-sterile. Unlike those one mycelia-sterile was identified as *Phaeosphaeria ammophilae* from rhizome by Eris *et al.*, [61]. One of the genus *Eurotium sp.* was also noted from the healthy rhizomes of *C. longa* [63].

3.2. Endophytic Fungi of Leaf

A total of 6 endophytic fungi have been isolated from leaves of *C. longa* through a study [64]. Another study carried by Eris *et al.*, have isolated one fungus *Daldinia eschscholzii* from leaf of *C. longa* [61].

Table 3. Endophytic fungi from different parts of turmeric plant.

Genus	Identified by	Plant tissue	References
<i>Acremonium sp.</i>	ITS-5.8s r-DNA analysis	Rh	[59]
<i>Aspergillus versicolor</i> and <i>Aspergillus venenatus</i>	ITS-5.8s r-DNA analysis	Rh	[59]
<i>Exophiala lecanii</i>	ITS-5.8s r-DNA analysis	Rh	[59]
<i>Fusarium Solani</i> , <i>Fusarium proliferatum</i> , <i>Fusarium oxysporum</i>	ITS-5.8s r-DNA analysis	Rh, S, Fl, In	[59] [60] [61]
<i>Trichoderma harzianum</i> , <i>Trichoderma atroviride</i> , <i>Trichoderma asperellum</i>	ITS-5.8s r-DNA analysis	Rh	[60]
<i>Penicillium citrinum</i>	ITS-5.8s r-DNA analysis	Rh	[59]
<i>Phoma herbarum</i>	ITS-5.8s r-DNA analysis	Rh	[59]
<i>Stachybotrys sp.</i>	ITS-5.8s r-DNA analysis	Rh	[59]
<i>Pestalopsis microspora</i>	ITS-5.8s r-DNA analysis	Rh	[60]
<i>Phaeosphaeria ammophilae</i>	ITS-5.8s r-DNA analysis	Rh	[61]
<i>Daldinia eschscholzii</i>	ITS-5.8s r-DNA analysis	L	[61]
<i>Arthrobotrys foliicola</i>	ITS-5.8s r-DNA analysis	R	[61]
<i>Drechslera kusanoi</i>	ITS-5.8s r-DNA analysis	In	[61]

Rh = Rhizome, S = Stem, Fl = Flower, In = Inflorescence, L = Leaf, R = Root.

3.3. Endophytic Fungi of Stem

Several endophytic fungi from fresh stem of *C. longa* have been isolated [62]. The endophytic fungus *Fusarium proliferatum* was isolated from stems of healthy *C. longa* collected in Indonesia [61].

3.4. Endophytic Fungi of Root

The endophytic fungus *Arthrobotrys foliicola* was isolated from the roots of *C. longa* [61]. Some endophytic fungi have also been isolated from roots of *C. longa* [62].

3.5. Endophytic Fungi of other Parts

Some endophytic fungi from tubers and young tubers of *C. longa* have been isolated [62]. Eris *et al.*, have isolated *Fusarium verticillioides* and *Phanerochaete chrysosporium* from the flowers of *C. longa* of which the later fungus is reported as mycelia-sterile. The group also isolated *Cochliobolus kusanoi* and *Fusarium proliferatum* from the inflorescence part of the plant. *Cochliobolus kusanoi* is the teleomorph stage of *Drechslera kusanoi* [61].

4. Reported Bioactivity on Endophytic Fungi Isolated from Turmeric Plant

4.1. Antioxidant Activity

44 endophytic fungi from several parts of turmeric plant were investigated by a research group Bustanussalam *et al.*, for their antioxidant bioactive compounds. They observed both filtrate and biomass ethylacetate extracts of endophytic fungi have antioxidant activity. On the basis of DPPH free radical scavenging method filtrate extracts gave higher antioxidant activity than biomass extracts. Due to presence of fatty acids, biomass extracts are less active as antioxidant. Filtrate extracts contain many secondary metabolites that may act as protection for the endophytic fungi of host plant [62].


4.2. Anticancer Activity

A study carried out by Ratih *et al.*, have shown inhibitory activity of 4 isolates of endophytic fungi towards human breast cancer MCF-7 cell line [65]. Endophytic fungus *Eurotium* sp. was isolated from rhizomes of *C. longa* by Ruby and Raghunath and it is utilized for the production of anticancer enzyme asparaginase for the first time [63]. The enzyme is considered to participate in major role in the asparaginase metabolism of the cells. It's a likely therapeutic agent for acute lymphocytic leukemia, acute lymphoblastic leukemia and chronic myelogenous leukemia also reported to be an effective antilymphoma agent in humans.

4.3. Antifungal Activity

Suruchi *et al.*, have separated endophytic fungi from *C. longa* for antifungal activity. The group reported the active constituent Gentisyl alcohol which is an an-

ti-phytopathogenic molecule from fungal isolate *P. herbarum*. The purified compound showed antagonistic activity against *C. gloeosporioides* causing leaf spot of turmeric and could, therefore, be explored as a biocontrol agent for the host-specific phytopathogen [59]. Similarly, Vinayarani and Prakash have isolated *T. harzianum* from the rhizome of turmeric and they concluded that this can be exploited as a potential biocontrol agent for suppressing rhizome rot and leaf blight diseases in turmeric [60].



Structure of Gentisyl alcohol

4.4. Antibacterial Activity

Endophytic fungi could be a promising source for antibacterial bioactive agents. A study carried out by Sharmila *et al.*, found that the endophytic fungus *Chaetomium sp.* isolated from *C. longa* exhibited protruding antibacterial activity against three different human pathogenic bacteria such as *Proteus sp.*, *Pseudomonas sp.* and *Staphylococcus aureus* [66].

5. Conclusion

Till today, the chemical composition of *C. longa* has thoroughly been investigated and a number of different active substances have been identified, which exhibit a wide range of medicinal value. But, very few structures of bioactive compounds are characterized and identified from endophytic fungi of *C. longa*. Most of the investigations are restricted to the level of fungal identification and bioactivity assay. But, endophytic fungi, producer of a wide array of secondary bioactive metabolites with their peculiar potential compounds, namely, antioxidant, antibacterial, antifungal, anticancer, etc., are the ideal targets for further drug discovery. These leads may play a major role in the recovery of infectious, inflammatory and also certain kinds of cancer diseases. So, extensive research is necessary in the area of endophytic fungi isolation and characterization of the compounds from *C. longa*.

References

- [1] Priyadarsini, K.I. (2014) The Chemistry of Curcumin: From Extraction to Therapeutic Agent. *Molecules*, **19**, 20091-20112.
<https://doi.org/10.3390/molecules191220091>
- [2] Chattopadhyay, I., Kaushik, B., Uday, B. and Ranajit, K.B. (2004) Turmeric and Curcumin: Biological Actions and Medicinal Applications. *Indian Academy of Sciences*, **87**, 44-53.

- [3] Chandrasekaran, C.V., Sundarajan, K., Edwin, J.R., Gururaja, G.M., Mundkinajedu, D. and Agarwal, A. (2013) Immunostimulatory and Anti-Inflammatory Activities of *Curcuma longa* Extract and Its Polysaccharide Fraction. *Pharmacognosy Research*, **5**, 71-79. <https://doi.org/10.4103/0974-8490.110527>
- [4] Hamid, N., Najmeh, S., Mortaza, R., Samira, R., Maryam, S. and Mahmoud, R.-K. (2014) Turmeric: A Spice with Multifunctional Medicinal Properties. *Journal of HerbMed Pharmacology*, **3**, 5-8. <http://www.herbmedpharmacol.com>
- [5] Bak, J.P., Kim, J.B., Park, J.H., Yang, Y.J., Kim, I.S., Choung, E.S. and Kang, S.C. (2011) Screening and Compound Isolation from Natural Plants for Anti-Allergic Activity. *Journal of the Korean Society for Applied Biological Chemistry*, **54**, 367-375. <https://doi.org/10.3839/jksabc.2011.058>
- [6] Ho, J.N., Park, S.J., Choue, R. and Lee, J. (2013) Standardized Ethanol Extract of *Curcuma longa* L. Fermented by *Aspergillus oryzae* Promotes Lipolysis via Activation of cAMP-Dependent PKA in 3T3-L1 Adipocytes. *Journal of Food Biochemistry*, **37**, 595-603. <https://doi.org/10.1111/jfbc.12011>
- [7] Nahid, A. and Tariq, M.H. (2012) Current Nutraceuticals in the Management of Osteoarthritis: A Review. *Therapeutic Advances in Musculoskeletal Disease*, **4**, 181-207. <https://doi.org/10.1177/1759720X11436238>
- [8] Yan, W., Bowen, W.D., Hopson, R., Mathew, A.E. and Jacob, J.N. (2013) Biological Studies of Turmeric Oil, Part 2: Isolation and Characterization of Turmeric Oil Components; Cytotoxic Activity against Pancreatic Cancer Cells. *Natural Product Communications*, **8**, 811-814. <https://doi.org/10.1177/1934578X1300800633>
- [9] Fagbemi, J.F., Ugoji, E., Adenipekun, T. and Adelowotan, O. (2009) Evaluation of the Antimicrobial Properties of Unripe Banana (*Musa sapientum* L.), Lemon Grass (*Cymbopogon citratus* S.) and Turmeric (*Curcuma longa* L.) on Pathogens. *African Journal of Biotechnology*, **8**, 1176-1182. <http://www.academicjournals.org/AJB>
- [10] Teow, S.-Y., Liew, K., Ali, S.A., Khoo, A.S.-B. and Peh, S.-C. (2016) Antibacterial Action of Curcumin against *Staphylococcus aureus*: A Brief Review. *Journal of Tropical Medicine*, **2016**, Article ID: 2853045. <https://doi.org/10.1155/2016/2853045>
- [11] Suresh, K.G., Nayaka, H., Dharmesh, S.M. and Salimath, P.V. (2006) Free and Bound Phenolic Antioxidants in amla (*Embllica officinalis*) and Turmeric (*Curcuma longa*). *Journal of Food Composition and Analysis*, **19**, 395-476. <https://doi.org/10.1016/j.jfca.2005.12.015>
- [12] Motterlini, R., Foresti, R., Bassi, R. and Green, C.J. (2000) Curcumin, an Antioxidant and Anti-Inflammatory Agent, Induces Heme Oxygenase-1 and Protects Endothelial Cells against Oxidative Stress. *Free Radical Biology & Medicine*, **28**, 1303-1312. [https://doi.org/10.1016/S0891-5849\(00\)00294-X](https://doi.org/10.1016/S0891-5849(00)00294-X)
- [13] Suryanarayana, P., Satyanarayana, A., Balakrishna, N., Kumar, P.U. and Reddy, G.B. (2007) Effect of Turmeric and Curcumin on Oxidative Stress and Antioxidant Enzymes in Streptozotocin-Induced Diabetic Rat. *Medical Science Monitor*, **13**, 286-292.
- [14] Ahmad, M., Kamran, S.H. and Mobasher, A. (2014) Protective Effect of Crude *Curcuma longa* and Its Methanolic Extract in Alloxanized Rabbits. *Pakistan Journal of Pharmaceutical Sciences*, **27**, 121-128. <https://www.researchgate.net/publication/259490723>
- [15] Liu, F., Wang, Y., Burkhart, T.A., Gonzalez Penedo, M.F., Ma, S.D., Choi, W., Lim, H.W. and Lee, H.Y. (2014b) Effect of Balanced Low Pressure Drying of *Curcuma longa* Leaf on Skin Immune Activation Activities. *Biomedical Materials*, **24**,

2025-2039.

- [16] Singh, G., Singh, O.P. and Maurya, S. (2002) Chemical and Biocidal Investigations on Essential Oils of Some Indian Curcuma Species. *Progress in Crystal Growth and Characterization of Materials*, **45**, 75-81.
[https://doi.org/10.1016/S0960-8974\(02\)00030-X](https://doi.org/10.1016/S0960-8974(02)00030-X)
- [17] Kheeree, N., Sangvanich, P., Puthong, S. and Karnchanatat, A. (2010) Antifungal and Antiproliferative Activities of Lectin from the Rhizomes of *Curcuma amarissima* Roscoe. *Applied Biochemistry and Biotechnology*, **162**, 912-925.
<https://doi.org/10.1007/s12010-009-8804-8>
- [18] De Alcantara, G.F.T., Simoes-Neto, E., Da Cruz, G.M.P., Nobre, M.E.P., Neves, K.R.T., De Andrade, G.M., De Castro Brito, G.A. and De Varros Viana, G.S. (2017) Curcumin Reverses Neurochemical, Histological and Immunohistochemical Alterations in the Model of Global Brain Ischemia. *Journal of Traditional and Complementary Medicine*, **7**, 14-23. <http://www.elsevier.com/locate/jtcm>
<https://doi.org/10.1016/j.jtcm.2015.10.001>
- [19] Ramos, A., Visozo, A., Piloto, J., García, A., Rodríguez, C.A. and Rivero, R. (2003) Screening of Antimutagenicity via Antioxidant Activity in Cuban Medicinal Plants. *Journal of Ethnopharmacology*, **87**, 241-246.
[https://doi.org/10.1016/S0378-8741\(03\)00156-9](https://doi.org/10.1016/S0378-8741(03)00156-9)
- [20] Hayashi, A. (2014) Effect on Lipid Metabolism by Turmeric (*Curcuma longa* Linn.) Supplementation in High Fructose-Diet-Fed Rats. *Bulletin of Tokyo Kasei University. 2, Natural Science*, **54**, 13-18.
- [21] Wang, F.W., Jiao, R.H., Cheng, A.B., Tan, S.H. and Song, Y.C. (2007) Antimicrobial Potentials of Endophytic Fungi Residing in *Quercus variabilis* and Brefeldin A Obtained from *Cladosporium* sp. *World Journal of Microbiology and Biotechnology*, **23**, 79-83. <https://doi.org/10.1007/s11274-006-9195-4>
- [22] Aggarwal, B.B., Sundaram, C., Malani, N. and Ichikawa, H. (2007) Curcumin: The Indian Solid Gold. *Advances in Experimental Medicine and Biology*, **595**, 1-75.
https://doi.org/10.1007/978-0-387-46401-5_1
- [23] Maheshwari, R.K., Singh, A.K., Gaddipati, J. and Srimal, R.C. (2006) Multiple Biological Activities of Curcumin: A Short Review. *Life Sciences*, **78**, 2081-2087.
<https://doi.org/10.1016/j.lfs.2005.12.007>
- [24] Chavalittumrong, P. and Jirawattanapong, W. (1992) Variation of Active Constituents of *Curcuma domestica* Rhizomes at Different Ages. *Thai Journal of Pharmaceutical Sciences*, **16**, 165-174.
- [25] Aggarwal, B.B., Kunnumakkara, A.B., Harikumar, K.B., Tharakan, S.T., Sung, B. and Anand, P. (2008) Potential of Spice-Derived Phytochemicals for Cancer Prevention. *Planta Medica*, **74**, 1560-1569. <https://doi.org/10.1055/s-2008-1074578>
- [26] Funk, J.L., Frye, J.B., Oyarzo, J.N., Kuscuoglu, N., Wilson, J., McCaffrey, G., Stafford, G., Chen, G., Lantz, R.C., Jolad, S.D., Sólyom, A.M., Kiela, P.R. and Timmermann, B.N. (2006) Efficacy and Mechanism of Action of Turmeric Supplements in the Treatment of Experimental Arthritis. *Arthritis & Rheumatology*, **54**, 3452-3464.
<https://doi.org/10.1002/art.22180>
- [27] Aggarwal, B.B., Bhatt, I.D., Ichikawa, H., Ahn, K.S., Sethi, G., Sandur, S.K., Sundaram, C., Seeram, N. and Shishodia, S. (2007) Curcumin-Biological and Medicinal Properties. CRC Press, Boca Raton, London, New York, 297-368.
- [28] Park, B.S., Kim, G.J., Kim, M.R., Lee, S.E., Takeoka, G.R., Oh, K.B. and Kim, J.H. (2005) *Curcuma longa* L. Constituents Inhibit Sortase A and *Staphylococcus aureus* Cell Adhesion to Fibronectin. *Journal of Agricultural and Food Chemistry*, **53**,

- 9005-9009. <https://doi.org/10.1021/jf051765z>
- [29] Roth, G.N., Chandra, A. and Nair, N.G. (1998) Novel Bioactivities of *Curcuma longa* Constituents. *Journal of Natural Products*, **61**, 542-545. <https://doi.org/10.1021/np970459f>
- [30] Li, W., Wang, S.S., Feng, J.T., Xiao, Y.S., Xue, X.Y., Zhang, H., Wang, Y.Q. and Liang, X.M. (2009) Structure Elucidation and NMR Assignments for Curcuminoids from the Rhizomes of *Curcuma longa*. *Magnetic Resonance in Chemistry*, **47**, 902-908. <https://doi.org/10.1002/mrc.2478>
- [31] Kita, T., Imai, S., Sawada, H. and Seto, H. (2009) Isolation of Dihydrocurcuminoids from Cell Clumps and Their Distribution in Various Parts of Turmeric (*Curcuma longa*). *Bioscience, Biotechnology, and Biochemistry*, **73**, 1113-1117. <https://doi.org/10.1271/bbb.80871>
- [32] Park, S.Y. and Kim, D.S.L.H. (2002) Discovery of Natural Products from *Curcuma longa* That Protect Cells from Beta-Amyloid Insult: A Drug Discovery Effort against Alzheimer's Disease. *Journal of Natural Products*, **65**, 1227-1231. <https://doi.org/10.1021/np010039x>
- [33] Kiuchi, F., Goto, Y., Sugimoto, N., Akao, N., Kondo, K. and Tsuda, Y. (1993) Nematocidal Activity of Turmeric: Synergistic Action of Curcuminoids. *Chemical and Pharmaceutical Bulletin*, **41**, 1640-1643. <https://doi.org/10.1248/cpb.41.1640>
- [34] Wang, L.Y., Zhang, M., Zhang, C.F. and Wang, Z.T. (2008) Diaryl Derivatives from the Root Tuber of *Curcuma longa*. *Biochemical Systematics and Ecology*, **36**, 476-480. <https://doi.org/10.1016/j.bse.2007.12.005>
- [35] Chen, J.J., Tsai, C.S., Hwang, T.L., Shieh, P.C., Chen, J.F. and Sung, P.J. (2010) Sesquiterpenes from the Rhizome of *Curcuma longa* with Inhibitory Activity on Superoxide Generation and Elastase Release by Neutrophils. *Food Chemistry*, **119**, 974-980. <https://doi.org/10.1016/j.foodchem.2009.07.060>
- [36] Zeng, Y.C., Qiu, F., Takahashi, K., Liang, J.M., Qu, G.X. and Yao, X.S. (2007) New Sesquiterpenes and Calebin Derivatives from *Curcuma longa*. *Chemical and Pharmaceutical Bulletin*, **55**, 940-943. <https://doi.org/10.1248/cpb.55.940>
- [37] Leela, N.K., Tava, A., Shafi, P.M., John, S.P. and Chempakam, B. (2002) Chemical Composition of Essential Oils of Turmeric (*Curcuma longa* L.). *Acta Pharmaceutica*, **52**, 137-141.
- [38] Chowdhury, J.U., Nandi, N.C., Bhuiyan, M.N. and Mobarok, M.H. (2008) Essential Oil Constituents of the Rhizomes of Two Types of *Curcuma longa* of Bangladesh. *Bangladesh Journal of Scientific and Industrial Research*, **43**, 259-266. <https://doi.org/10.3329/bjsir.v43i2.970>
- [39] Gopalan, B., Goto, M., Kodama, A. and Hirose, T. (2000) Supercritical Carbon Dioxide Extraction of Turmeric (*Curcuma longa*). *Journal of Agricultural and Food Chemistry*, **48**, 2189-2192. <https://doi.org/10.1021/jf9908594>
- [40] Usman, L.A., Hamid, A.A., George, O.C., Ameen, O.M., Muhammad, N.O. and Zubair, M.F.A.L. (2009) Chemical Composition of Rhizome Essential Oil of *Curcuma longa* L. Growing in North Central Nigeria. *World Journal of Chemistry*, **4**, 178-181.
- [41] Ma, X. and Gang, D.R. (2006) Metabolic Profiling of Turmeric (*Curcuma longa* L.) Plants Derived from *in Vitro* Micropropagation and Conventional Greenhouse Cultivation. *Journal of Agricultural and Food Chemistry*, **54**, 9573-9583. <https://doi.org/10.1021/jf061658k>
- [42] Zeng, Y.C., Liang, J.M., Qu, G.X. and Qiu, F. (2007) Chemical Constituents of *Cur-*

- cuma longa* I: Bisabolane Sesquiterpenes. *Acta Pharmaceutica Sinica*, **17**, 738-741.
- [43] Chassagnez-Me'ndez, A.L., Machado, N.T., Araujo, M.E., Maia, J.G. and Meireles, M.A.A. (2000) Supercritical CO₂ Extraction of Curcumins and Essential Oil from the Rhizomes of Turmeric (*Curcuma longa* L.). *Industrial & Engineering Chemistry Research*, **39**, 4729-4733. <https://doi.org/10.1021/ie000171c>
 - [44] Li, W., Feng, J.T., Xiao, Y.S., Wang, Y.Q., Xue, X.Y. and Liang, X.M. (2009) Three Novel Terpenoids from the Rhizomes of *Curcuma longa*. *Journal of Asian Natural Products Research*, **11**, 569-575. <https://doi.org/10.1080/10286020902939182>
 - [45] Golding, B.T., Pombo, E. and Christopher, J.S. (1982) Turmerones: Isolation from Turmeric and Their Structure Determination. *Journal of the Chemical Society, Chemical Communications*, **6**, 363-364. <https://doi.org/10.1039/c39820000363>
 - [46] Imai, S., Morikiyo, M., Furihata, K., Hayakawa, Y. and Seto, H. (1990) Turmeronol A and Turmeronol B, New Inhibitors of Soybean Lipoxygenase. *Agricultural and Biological Chemistry*, **54**, 2367-2371. <https://doi.org/10.1080/00021369.1990.10870290>
 - [47] Manzan, A.C.C.M., Toniolo, F.S., Bredow, E. and Povh, N.P. (2003) Extraction of Essential Oil and Pigments from *Curcuma longa* [L.] by Steam Distillation and Extraction with Volatile Solvents. *Journal of Agricultural and Food Chemistry*, **51**, 6802-6807. <https://doi.org/10.1021/jf030161x>
 - [48] Nishiyama, T., Mae, T., Kishida, H., Tsukagawa, M., Mimaki, Y., Kuroda, M., Sashida, Y., Takahashi, K., Kawada, T., Nakagawa, K. and Kitahara, M. (2005) Curcuminoids and Sesquiterpenoids in Turmeric (*Curcuma longa* L.) Suppress an Increase in Blood Glucose Level in Type 2 Diabetic KK-Ay Mice. *Journal of Agricultural and Food Chemistry*, **53**, 959-963. <https://doi.org/10.1021/jf0483873>
 - [49] Ohshiro, M., Kuroyanagi, M. and Ueno, A. (1990) Structures of Sesquiterpenes from *Curcuma longa*. *Phytochemistry*, **29**, 2201-2205. [https://doi.org/10.1016/0031-9422\(90\)83038-3](https://doi.org/10.1016/0031-9422(90)83038-3)
 - [50] Wang, L.Y., Zhang, M., Zhang, C.F. and Wang, Z.T. (2008) Alkaloid and Sesquiterpenes from the Root Tuber of *Curcuma longa*. *Acta Pharmaceutica Sinica*, **43**, 724-727.
 - [51] Braga, M.E., Leal, P.F., Carvalho, J.E. and Meireles, M.A. (2003) Comparison of Yield, Composition, and Antioxidant Activity of Turmeric (*Curcuma longa* L.) Extracts Obtained Using Various Techniques. *Journal of Agricultural and Food Chemistry*, **51**, 6604-6611. <https://doi.org/10.1021/jf0345550>
 - [52] Awasthi, P.K. and Dixit, S.C. (2009) Chemical Composition of *Curcuma longa* Leaves and Rhizome Oil from the Plains of Northern India. *Pharmacognosy*, **1**, 312-316. <https://doi.org/10.4103/0975-1483.59319>
 - [53] Liu, C., Sun, B., Huang, J., Gao, H., Wen, S. and Wu, L. (2007) A Novel Sesquiterpene from *Curcuma longa*. *Asina Journal of Traditional Medicines*, **2**, 82-84.
 - [54] Mohamed, S.M., El-Gengaihi, S.E. and Motawe, H.M. (2003) Terpenoid from *Curcuma longa*. *Egyptian Journal of Pharmaceutical Sciences*, **43**, 139-151.
 - [55] Wu, Z.H., Huang, S.W., Liu, C.Y., Sun, B.H., Huang, J.S. and Wu, L.J. (2008) A Novel Compound from *Curcuma longa*. *Asina Journal of Traditional Medicines*, **3**, 199-202.
 - [56] Ashraf, A. and Javaid, A. (2005) Fungi Associated with Rhizome of Turmeric (*Curcuma longa* L.) in Pakistan. *Mycopath*, **3**, 69-71.
 - [57] Singh, D., Rathod, V., Ninganagouda, S., Herimath, J. and Kulkarni, P. (2013) Biosynthesis of the Silver Nanoparticle by Endophytic Fungi *Penicillium* sp. Isolated

- from *Curcuma longa* (Turmeric) and Its Antibacterial Activity against Pathogenic Gram-Negative Bacteria. *Journal of Pharmacy Research*, **7**, 448-453.
<https://doi.org/10.1016/j.jopr.2013.06.003>
- [58] Maehara, S., Ikeda, M., Haraguchi, H., Kitamura, C., Nagoe, T., Ohashi, K. and Shibuya, H. (2011) Microbial Conversion of Curcumin into Colorless Hydro Derivatives by the Endophytic Fungus *Diaporthe* sp. Associated with *Curcuma longa*. *Chemical and Pharmaceutical Bulletin*, **59**, 1042-1044.
<https://doi.org/10.1248/cpb.59.1042>
- [59] Suruchi, G., Sanjana, K., Baljinder, S., Ram, A.V. and Manoj, K.D. (2016) Production of Gentisyl Alcohol from *Phoma herbarum* Endophytic in *Curcuma longa* L. and Its Antagonistic Activity towards Leaf Spot Pathogen *Colletotrichum gloeosporioides*. *Applied Biochemistry and Biotechnology*, **180**, 1093-1109.
<https://doi.org/10.1007/s12010-016-2154-0>
- [60] Vinayarani, G. and Prakash, H.S. (2018) Fungal Endophytes of Turmeric (*Curcuma longa* L.) and Their Biocontrol Potential against Pathogens *Pythium aphanidermatum* and *Rhizoctonia solani*. *World Journal of Microbiology and Biotechnology*, **34**, 49. <https://doi.org/10.1007/s11274-018-2431-x>
- [61] Eris, S., Nampiah, S. and Partomuan, S. (2017) Endophytic Fungi Associated with Turmeric (*Curcuma longa* L.) Can Inhibit Histamine-Forming Bacteria in Fish. *HAYATI Journal of Biosciences*, **24**, 46-52.
<https://doi.org/10.1016/j.hjb.2017.05.004>
- [62] Bustanussalam, R.F., Septiana, E., Lekatompeppy, S.J., Widowati, T., Sukiman, H.I. and Simanjuntak, P. (2015) Screening for Endophytic Fungi from Turmeric Plant (*Curcuma longa* L.) of Sukabumi and Cibinong with Potency as Antioxidant Compounds Producer. *Pakistan Journal of Biological Sciences*, **18**, 42-45.
<https://doi.org/10.3923/pjbs.2015.42.45>
- [63] Ruby, E.J. and Mahajan, R.T. (2014) Production of Anticancer Enzyme Asparaginase from Endophytic *Eurotium* Sp. Isolated from Rhizomes of *Curcuma longa*. *European Journal of Experimental Biology*, **4**, 36-43.
- [64] Dattu, S., Vandana, R., Ashish, K.S., Manzoor, U.H., Jasmine, M. and Prema, K. (2016) A Study on Extracellular Synthesis of Silver Nanoparticles from Endophytic Fungi, Isolated from Ethanomedicinal Plants *Curcuma longa* and *Catharanthus roseus*. *International Letters of Natural Sciences*, **57**, 58-66.
<https://doi.org/10.18052/www.scipress.com/ILNS.57.58>
- [65] Ratih, A.N., Bustanussalam, P.H.W., Neng, H., Adi, S. and Partomuan, S. (2017) Inhibitory Activity of Endophytic Fungi Isolated from Sukabumi Turmeric Plant (*Curcuma longa* L.) towards MCF-7 Cell Line. *Indonesian Journal of Pharmacy*, **28**, 19-25. <https://doi.org/10.14499/indonesianjpharm28iss1pp19>
- [66] Sharmila, T., Sreelatha, G.L., Geethanjali, P. and Subramanya, S. (2017) Antibacterial Activities of Endophytic Fungal Strains Isolated from Aromatic Plants. *Asian Journal of Pharmaceutical Science & Technology*, **7**, 28-33.